


Scottish Burden of Disease Study, 2015

Chronic liver diseases (including cirrhosis)
technical overview



This resource may also be made available on request in the following formats:



 0131 314 5300

 nhs.healthscotland-alternativeformats@nhs.net

Published by NHS Health Scotland

1 South Gyle Crescent
Edinburgh EH12 9EB

© NHS Health Scotland 2017 and ISD 2017

Background

The Scottish Burden of Disease (SBoD) study team have published comprehensive estimates of the burden of disease and injury in Scotland for 2015 [1]. The purpose of this technical overview is to provide background information on the data and methodology used, noting any caveats associated with estimating the burden of chronic liver disease, including cirrhosis (CLD) in SBoD.


Burden of disease studies aim to estimate the difference between ideal and actual health in a country or region at a specific point in time. Individuals can suffer non-fatal health loss due to suffering disability attributable to a disease, condition or injury, or suffer fatal health loss which is early death due to a disease, condition or injury. To quantify the total burden, non-fatal and fatal health loss are combined to produce a single metric called the Disability-Adjusted Life Year (DALY).

Further information about the SBoD study, including a more thorough explanation of the methodology used, overview reports, detailed results and other specific disease briefings, can be found on the website of the Scottish Public Health Observatory (ScotPHO) [1].

Estimated burden due to chronic liver disease

CLD was the 11th most common cause of disease burden in Scotland in 2015, resulting in a total of approximately 28,200 DALYs. Of this total burden, 87% was due to premature mortality attributed to CLD and 13% was attributed to health loss suffered due to living with CLD.

Figure 1 Percentage of total DALYs by gender and age-group for CLD



Men contributed a higher proportion of the burden (58%) than women (42%). Overall, 94% of the total CLD burden was contributed by individuals aged 35 years and over, as outlined in Figure 1. Note that the burden we are describing is the absolute burden and has not been adjusted for the age/gender case-mix.

How did we produce these estimates?

DALYs attributed to a disease, condition or injury are calculated by combining estimates from two individual metrics: Years of Life Lost (YLL) due to premature mortality and Years Lived with Disability (YLD).

Years of life lost (YLL) due to chronic liver disease

YLL measures the years of life lost due to premature deaths i.e. the fatal component of burden of disease. YLLs are calculated by subtracting the age at each CLD death from the expected remaining life expectancy for a person at that age.

Estimating the number of deaths

There were approximately 1,100 deaths caused by CLD in 2015. These deaths were identified from the underlying cause of death on the National Records of Scotland (NRS) register of deaths [2]. To classify deaths the GBD 2015 cause list was used,

which has been created using the International Statistical Classification of Diseases and Related Health Problems (ICD-10) [3, 4]. The NRS register of deaths has a Community Health Index (CHI) number attached to each death, which allows for demographic data such as gender, geographical area of residence and age at death to be established for each individual.

Included in the total CLD mortality count are deaths that have come from what are termed ill-defined causes of death in burden of disease studies. These ill-defined deaths are causes of death that have been coded with ICD-10 codes in vital registers but for the purposes of burden of disease studies, are not regarded as sufficiently specific causes of death. These ill-defined deaths are therefore redistributed amongst specific causes of death across the burden of disease cause list based on the redistribution of deaths method used in the GBD study [3]. For CLD, approximately 7% of the mortality count comes from ill-defined death categories such as 'unspecified haematemesis, melaena and gastrointestinal haemorrhage'. Further explanation of this method is available in the SBoD technical paper [1]. For this reason, the number of deaths due to CLD which have been reported are different from that of officially reported sources.

Life expectancy and YLL

Each single death contributes to the total YLL through calculating the difference between the age at death and the life expectancy at that age. Life expectancy was defined using the 2013 gender-specific National Life Tables for Scotland [5]. There were approximately 24,600 YLL due to CLD in Scotland in 2015. Dividing the total YLL for CLD by the total mortality count indicates that, on average, individuals who die due to CLD die approximately 22 years earlier than would otherwise be expected on the basis of the life expectancy of the general population.

Years lived with disability (YLD) due to chronic liver disease

Years lived with disability (YLD) are estimated using:

- disease and injury prevalence estimates
- levels of severity
- disability weights

Our sources of information for these three components are as follows:

Estimating the prevalence

To estimate prevalent cases of CLD in 2015, the Scottish Morbidity Records 01 (SMR01) was used [6]. This dataset contains structured data in the form ICD-10 codes relating to diagnoses made on discharge from general and acute hospitals during inpatient episodes and day cases. There are up to six individual ICD-10 codes that can be recorded, where the primary diagnosis relates to the main reason for the hospital episode of care, and the other secondary diagnoses refer to co-morbidities that may affect care during that hospital episode of care.

The SMR01 dataset has a CHI number attached to the hospital episode of care, which allows for the identification of records for an individual. This CHI number has been linked to records from the NRS register of deaths, to exclude individuals that have died from prevalence estimates that relate to a period following their date of death [2]. The number of individuals that had a primary diagnosis of CLD between 1 January 2006 to 31 December 2015 was used to estimate the number of prevalent cases.

Using this method of identifying prevalent cases of CLD, we estimated that there were approximately 21,800 individuals in the Scottish population living with disability due to CLD in 2015.

Severity distribution and disability weights

The levels of severity and disability due to CLD in Scotland were based on the specifications of the GBD 2015 study [7]. This allowed us to disaggregate the

prevalent cases into levels of severity and the associated disability at each level of severity. The disability weights were developed by the GBD study through surveys of the general public and take into account the consequences of each disease, condition and injury [8]. The severity distribution and disability weights for CLD are outlined in Table 1.

Table 1 Description and allocation to severity levels for CLD with corresponding disability weight

Severity level	Description	% of individuals	Disability weight
All	Has a swollen belly and swollen legs. The person feels weakness, fatigue and loss of appetite.	100	0.178

Once the severity of CLD and associated disability were taken into account, individuals were estimated to be suffering approximately 3,600 YLDs in 2015 due to living with CLD.

Data quality

In order to provide a measure of the degree of accuracy¹ and relevance² of the estimated disease DALYs to users, a measure of data quality has been developed for the SBoD study. This measure assigns a RAG (Red; Amber; Green) status to each disease or injury indicative of the accuracy and relevance of the estimates. Interpretation of the RAG status can be defined as follows:

Highly accurate and relevant

Estimates have been derived using relevant and robust data sources with only a small degree of adjustments performed to the input data. These estimates can be considered a highly accurate depiction of the burden incurred from the disease, condition or injury.

¹ How precise, unbiased or certain the estimate is.

² Do we measure the thing we want to measure?

 **Moderately accurate and relevant**

Estimates have been derived using reasonably relevant and robust data sources with only a moderate degree of adjustments performed to the input data. These estimates can be considered a moderately accurate depiction of the burden incurred from the disease, condition or injury.

 **Uncertainties over accuracy and relevance**

Estimates have been derived using less comprehensive or relevant data sources with a high degree of adjustments performed to the input data. These estimates contain substantial uncertainties and should be used with some caution.

The data quality has been assessed using three main criteria:

- Relevance and accuracy of the data source used to measuring the population of interest
- Likelihood that the implemented disease model captured the overall burden of disease or injury
- The relative contribution of ill-defined deaths to YLL, and YLL to DALY.

These criteria are subjectively assessed and each criterion is scored on a scale of 1 to 5. Further details on these data quality measures are available on the ScotPHO website [1].

Based on these criteria, the estimates of burden of CLD in Scotland are green,

 **highly accurate and relevant.**

We have chosen to use hospital diagnosis records to estimate the number of individuals suffering disability due to CLD. Due to the progression of CLD, which can result in health loss and poor outcomes, hospital records are likely to be a good resource to estimate prevalence, especially in those patients that have CLD for some time and whose liver is unable to compensate for prior cirrhotic damage [9].

Our study estimated a CLD prevalence of 0.41% in Scotland in 2015. In comparison, the Global Burden of Disease study (GBD) 2015 estimated a much lower CLD

prevalence of 0.05% [10]. A study carried out across the UK, using records from primary care over a 10-year period, estimated a prevalence of 0.76% for individuals aged 25 and above [11]. The same study also indicated that 38% of CLD was alcohol-related. Given the causal link between CLD and alcohol consumption, combined with the fact there are known issues with excess alcohol use in Scotland [12], we believe that using Scottish health records to describe the burden would be the most relevant approach.

What next to improve estimates for chronic liver disease?

Future work on the SBoD study will include reviewing the coding and recording of CLD in alternative national datasets. The development of the Scottish Primary Care Information Resource (SPIRE) will provide another source to estimate the prevalence of CLD for comparison [13]. Further to this, work will be carried out to attempt to derive estimates of severity levels that are dependent on age and that are specific to the Scottish population.

These improvements are partly dependant on exploring other data sources and reviewing evidence from high quality research that it is relevant to Scotland. Please contact the SBoD project team (nhs.healthscotland-sbod-team@nhs.net) for enquiries and suggestions on how to improve our estimates.

References

- [1] Scottish Burden of Disease study. Scottish Public Health Observatory, Available from: URL: <http://www.scotpho.org.uk/comparative-health/burden-of-disease/overview> (Accessed 03 July 2017)
- [2] National Records of Scotland (NRS). Vital Events - Deaths. Scottish Government, Available from: URL: <https://www.nrscotland.gov.uk/statistics-and-data/statistics/statistics-by-theme/vital-events/deaths> (Accessed 03 July 2017)
- [3] GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet* 2016 Oct 8;388(10053):1459-544.
- [4] World Health Organization. International Statistical Classification of Diseases and Related Health Problems 10th Revision (ICD-10) Version for 2010, Geneva. World Health Organization, Available from: URL: <http://apps.who.int/classifications/icd10/browse/2010/en> (Accessed 03 July 2017)
- [5] Office for National Statistics. National Life Tables: Scotland. Open Government Licence, Available from: URL: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/lifeexpectancies/datasets/nationallifetables/scotlandreferencetables> (Accessed 03 July 2017)
- [6] ISD Scotland. SMR Datasets - Structuring Healthcare Information, <http://www.isdscotland.org/products-and-services/structuring-healthcare-information/CHH-IST/SMRDS-HTML2/index.html> (Accessed 03 July 2017)

- [7] Vos T, Allen C, Arora M, Barber RM, Bhutta ZA, Brown A, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet* 2016;388(10053):1545-602.
- [8] Salomon JA, Haagsma JA, Davis A, de Noordhout CM, Polinder S, Havelaar AH, et al. Disability weights for the Global Burden of Disease 2013 study. *The Lancet Global Health* 2015 Nov;3(11):e712-e723.
- [9] Volk ML, Tocco RS, Bazick J, Rakoski MO, Lok AS. Hospital readmissions among patients with decompensated cirrhosis. *Am J Gastroenterol* 2012;107:247-252.
- [10] Institute for Health Metrics and Evaluation (IHME). GBD Results Tool. Seattle, WA: IHME, University of Washington. Available from: URL: <http://ghdx.healthdata.org/gbd-results-tool> (Accessed 03 July 2017)
- [11] Fleming KM, Aithal GP, Solaymani-Dodaran M, Card TR and West J. Incidence and prevalence of cirrhosis in the United Kingdom, 1992–2001: A general population-based study. *Journal of Hepatology* 2009 Jul;51(1):233-234.
- [12] Beeston C, Reid G, Robinson M, Craig N, McCartney G, Graham L and Grant I (on behalf of the MESAS project team). *Monitoring and Evaluating Scotland's Alcohol Strategy. Third Annual Report*. Edinburgh: NHS Health Scotland; 2013
- [13] Scottish Primary Care Information Resource (SPIRE). NHS National Services Scotland, Available from: URL: <http://spire.scot/> (Accessed 03 July 2017)

